

Product Datasheet

Name: Mouse Anti-SARS-CoV-2 N protein Monoclonal Antibody

Description: Hybridoma clones have been derived from hybridization of myeloma cells with spleen cells of BALB/c mouse immunized with recombinant SARS-CoV-2 N protein.

| Catalog No. | lsotype | Clone No. | Usage | Buffer |
|-------------|---------|-----------|--------------------|------------------|
| bsm-41411M | lgG2b | 1C7 | Capture /Detection | 10mM PBS (pH7.4) |

Specificity: Mab react with recombinant antigen SARS-CoV-2 N protein

Host: Mouse

Clonality: Monoclonal

Format: Liquid

Concentration: ≥1 mg/ml

Purification: ≥90% (SDS-PAGE)

Preservative: 0.1% Proclin300

Application: Recommended for sandwich immunoassays in ELISA and CLIA. Each laboratory should determine an optimum working titer for use in its particular application.

Storage: Store at -20 °C for three years. Avoid repeated freeze/thaw cycles.

Background: Nucleocapsid protein is a most abundant protein of coronavirus. During virion assembly, N protein binds to viral RNA and leads to formation of the helical nucleocapsid. Nucleocapsid protein is a highly immunogenic phosphoprotein also implicated in viral genome replication and in modulating cell signaling pathways. Because of the conservation of N protein sequence and its strong immunogenicity, the N protein of coronavirus is chosen as a diagnostic tool.


Note: This product as supplied is intended for research or further manufacturing use only.



400-901-9800
sales@bioss.com.cn
support@bioss.com.cn

VALIDATION IMAGES

kDa M A B C D E F



Anti-SARS-CoV-2 N protein Mouse Monoclonal Antibody (Cat# bsm-41411M) at 1:1000 dilution. Sample: rSARS-CoV-2 N Protein (Cat# bs-41408P)

Lane A: 50ng

Lane B: 25ng

Lane C: 5ng

Lane D: 2ng

Lane E: 1ng

Lane F: 0.5ng

Secondary

Goat Anti-Mouse IgG (H+L)/IRDye800CW at 1/20000 dilution.

Developed using the Odyssey (Li-cor).

Performed under reducing conditions.